

Breastfeeding and Consumption Food

by I Rai Ngardita

Submission date: 20-Apr-2023 03:02PM (UTC+0700)

Submission ID: 2070148072

File name: The_Influence_of_Factors_of_Breastfeeding__19.pdf (2.12M)

Word count: 1024

Character count: 5558

The Influence of Factors of Breastfeeding and Food Consumption on Stunting Incidence among Preschool Children in Keerom District, Papua Province, Indonesia

By Sanya Lusiana

WORD COUNT

5412

TIME SUBMITTED

09-MAR-2022 04:41PM

PAPER ID

83444801



The Influence of Factors of Breastfeeding and Food Consumption on Stunting Incidence among Preschool Children in Keerom District, Papua Province, Indonesia

I. Rai Ngardita^{1*}, Ratih Nurani Sumardi², Sanya Anda Lusiana³, I Komang Agusjaya Mataram², Andi Eka Yunianto³

¹Department of Nutrition, Health Polytechnic of Health-Ministry, Jayapura, Indonesia; ²Department of Nutrition, Health Polytechnic of Health-Ministry, Denpasar, Indonesia; ³Department of Nutrition, Faculty of Health Science, Siliwangi University, Tasikmalaya, Indonesia

Abstract

Citation: Ngardita I, Sumardi RN, Lusiana SA, Mataram KA, Yunianto AE. The Influence of Factors of Breastfeeding and Food Consumption on Stunting Incidence among Preschool Children in Keerom District, Papua Province, Indonesia. Open Access Maced J Med Sci. 2021 Jul 07; 9(E):559-565. <https://doi.org/10.3889/oamjms.2021.6076>

Keywords: Breastfeeding; Consumption; Stunting
***Correspondence:** Ngardita, Department of Nutrition, Health Polytechnic of Health-Ministry, Jayapura, Indonesia. E-mail: raingardita@gmail.com

Received: 23-Mar-2021
Revised: 30-May-2021
Accepted: 17-Jun-2021

Copyright: © 2021 I. Rai Ngardita, Ratih Nurani Sumardi, Sanya Anda Lusiana, I Komang Agusjaya Mataram, Andi Eka Yunianto

Funding: This research did not receive any financial support

Competing Interests: The authors have declared that no competing interests exist

Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: This study was a nutritional problem due to linear growth failure in children under five, resulting in future health problems. Papua has a prevalence of stunting in children under five of 17.8% in 2018.

AIM: This study aimed to analyze the influence of breastfeeding and consumption factors on the incidence of stunting in Keerom District, Papua Province.

METHOD: This study was a cross-sectional design. The sample in this study was 324 toddlers. Characteristics data and breastfeeding were obtained by interviewing parents. In contrast, data on the consumption of children under five were obtained using the Food Recall. Nutritional status was calculated using the Z-score obtained from measurements of body weight and height of children under five. SEM analysis was used to analyze breastfeeding and consumption factors on children's nutritional status under five.

RESULT: The study results based on consumption ($p < 0.05$; $r = 0.052$) and breastfeeding ($p < 0.05$; $r = -0.015$) had a relationship with the incidence of stunting.

CONCLUSION: There was a significant effect of consumption and breastfeeding factors on the incidence of under-five stunting.

Introduction

The first global nutrition goal for 2030 by the World Health Assembly is to reduce the proportion of children under five who stunted by 40% [1]. Stunting reflects the condition of failure to thrive in children under five due to chronic malnutrition. It is characterized by a height or weight that is not sufficient for the growth reference standard, especially at 3–5 years [2]. This linear growth failure is a marker of various pathological disorders associated with increased morbidity and mortality, loss of physical growth potential, decreased neurodevelopmental and cognitive function, and an increased risk of chronic disease in adulthood [3].

Chronic malnutrition occurs when the baby is in womb, but stunting will be seen after the baby is 2 years old. Therefore, the first 1000 days of life become the main focus because it determines the level of physical growth, intelligence, and productivity in the future. The 2013 Riskesdas results show that around 37% (9 million) children are stunted,

and Indonesia ranks fifth with the world's highest stunting prevalence. Toddlers and Baduta who are stunted will be more susceptible to disease and are at risk of decreasing productivity levels [4]. If this situation is not resolved, it can affect Indonesia's development performance, economic growth, poverty, and inequality. In Indonesia, stunted children are experienced by poor or underprivileged families and non-poor families.

According to WHO, the causes of stunting in children are divided into two factors: direct and indirect causes. The immediate causes are family factors (pregnancy and environment), inadequate feeding (poor quality of food, wrong feeding, and sanitation), breastfeeding practices, and infections. Food consumption is one of the causes of stunting in children under five. Toddlers need quality nutrient-rich foods that are obtained by eating a variety of foods [5], [6], [7]. Poor diet results in malnutrition in children, further exacerbated by micronutrient deficiencies, which can worsen the nutritional status and level of health of children [8]. Besides, breastfeeding has been identified

as a modifiable health behavior to protect children from stunting [9], [10]. Lack of consumption of breast milk can have an effect on increasing the risk of growth and development in preschool children [11].

Proper breastfeeding is the most effective step to meet children's nutritional adequacy. The nutritional composition in breast milk fulfills the child's dietary needs and positively influences children's growth [12]. Early initiation of breastfeeding may prevent and reduces the risk of stunting, while delaying breastfeeding initiation resulted in a 1.3 times higher risk of stunting [8]. It is hoped that breastfeeding for a minimum of 6 months or exclusive breastfeeding is also essential to reduce stunting prevalence in children under five [9].

Papua Province is one of the provinces that has a higher stunting rate than the national one. Riskesdas 2018 show the prevalence of nutritional status in children aged 0–23 months (Baduta) is very short at 15.1%, while in Indonesia, it is 12.8%. In Papua, it was 18.9%, and Indonesia 17.1%. The prevalence of nutritional status in very short children aged 0–59 months (toddlers) is 15.3%, while in Indonesia, it is 11.5%. The prevalence of stunting is 17.8%; this figure is lower than in Indonesia, which is 19.3% [13].

In 2018 the National Team for the Acceleration of Poverty Reduction (TNP2K), the Coordinating Ministry for Human Development and Culture, and Bapenas set priority districts/cities for handling stunting in Indonesia. In 2019 (phase II of handling stunting), the priority districts/towns in Papua Province are Asmat, Biak Numfor, Boven Digoel, Deyai, Keerom, and Yapen Islands [14]. Keerom District is one of the districts in Papua Province, located south of Jayapura City, and can be reached by land route. The prevalence of stunting in the Keerom District is 10.45%, with a poverty rate of 45.11%. Keerom District consists of 7 districts and 61 villages. Keerom District is one of the districts included in phase II in handling stunting in the Papua region.

Based on the description above, it is necessary to analyze the risk factors associated with stunting problems, especially in children under five in Keerom District. These factors are then analyzed statistically using structural equation modeling (SEM) to assess the strength of their relationship with one another. The results of this SEM analysis will be used to formulate a model for the occurrence of nutritional problems in children under five, which link stunting with family and household factors. Complementary feeding is inadequate in breastfeeding practices and infectious diseases. With this model's compilation, it is hoped that policyholders at the Keerom District level, such as the Health Office, will focus more on developing specific nutrition problem management programs based on risk factor analysis in each working area of Keerom District.

Methods

This research received ethical approval from the health research ethics committee of the Health Polytechnic of the Ministry of Health Yogyakarta with the number: e-KEPK/POLKESYO/0371/X/2019. This research was conducted for 15 months, starting from August to December 2019. This type of research was a cross-sectional design. The study was located in Keerom District, Papua Province, one of the 160 priority regions for stunting intervention with a prevalence of 10.45%. The sampling technique was carried out by stratified random sampling. Of the 9 stunting locus areas in the Keerom region, three regions were randomly selected, then 30% of the under-five population in the three regions were selected again. About 324 children under five were involved as samples. The inclusion criteria in this study including (a) children aged 6–59 months, (b) having a biological mother, (c) parents willing to participate in the study. The criteria for children under five, including (a) not sick or on outpatient treatment, (b) not at the research site at the time of the study, (c) under-fives with congenital disorders. The exclusion criteria were children under five who were sick or on outpatient treatment/not at the research location at the time of the study, (b) under-fives with congenital disorders (c) staying at the research location for <6 months. The variables examined in this study were nutritional status as the dependent variable and breastfeeding and consumption as independent variables.

Socio-demographic characteristics

The socio-geographical characteristics data in this study consisted of toddler characteristics and parental characteristics. The data on children under five's characteristics gender, age, height, and parental characteristics included age, education, and occupation.

Breastfeeding practices

Breastfeeding data in this study consisted of early breastfeeding initiation practices, exclusive breastfeeding, and duration of breastfeeding.

Consumption

Consumption data were collected using the 1x24 hour food recall method for two days. This study's consumption data consisted of energy, protein, Vitamin A, iron, iodine, and zinc. Nutritional adequacy is calculated directly by comparing the nutritional adequacy rate. Furthermore, the level of nutritional adequacy is obtained by comparing the amount of consumption of these nutrients with their adequacy [15]. The following is the formula for the adequacy of nutrients used:

$TKGi = (Ki/AKGi) \times 100\%$

Note:

$TKGi$ = The level of adequacy of the nutrient i

Ki = Consumption of nutrients i

$AKGi$ = Adequacy rate of the nutrient i .

Nutritional status

The nutritional status of children under five is determined based on the WHO Growth Child standards 2006 and calculated by the WHO Anthro software (version 3.12, January 2011). Toddler status data includes: height-for-age z-score (HAZ), Weight-for-age z-score (WAZ), Weight-for-height z-score (WHZ), and BMI-for-age z-score (zBMI). The following cut-offs as defined by the WHO were used: stunted: <-2 HAZ (moderate stunted: $-3 \leq HAZ <-2$; severe stunted: $HAZ <-3$); acute malnutrition based on WHZ score: <-2 WHZ (acute malnutrition: $-3 \leq WHZ <-2$; severe acute malnutrition: $WHZ <-3$).

Data analysis

The data that have been collected was processed according to the type of variable. The process of entering data (data entry) using a computer. To determine the strength of the relationship between variables and to compile a stunting model in Keerom District, the AMOS 20 structural equation modeling (SEM) analysis was used.

Nutritional status indicators were body weight for age (NS1), height for age (NS2), weight for height (NS3), and body mass index for age (NS4). Indicators of breastfeeding were early breastfeeding (BF1), exclusive breastfeeding (BF2), and duration of breastfeeding (BF3). Consumption indicators were energy consumption (CF1), protein (CF2), iron (Fe) (CF3), Vitamin A (CF4), zinc (Zn) (CF5), Iodine (CF6), meal frequency (CF7), meal arrangement (CF8), and availability of foodstuffs (CF9).

Results

This research was conducted in Keerom Papua District. Specifically, the study was located in the Arso Kota and Arso III Community health centers (Puskesmas) areas. It involving five villages, namely Arso Kota, Jaifuri, Gudang Garang, Alang-Alang, and the sample in this study was 324 samples. Respondents in this study were mothers of toddlers. This study's sample was toddlers with no sick criteria and was in the study location. The distribution of samples according to the characteristics can be seen in the following table.

Table 1 shows that the distribution of male and female respondents was almost the same. The sample's mean age was 27.77 months (2 years three months), and the mean sample weight at birth was 3.30 kg.

Table 1: Distribution of samples according to characteristics

Characteristics of the sample	f	%
Gender		
Male	165	50.9
Female	159	49.1
Age (months) (mean \pm SD)	27.77 \pm 14.82	
Birth weight (kg) (mean \pm SD)	3.30 \pm 0.46	

Table 2 shows that the father's mean age was 34 years and the mother's average age was 27 years. Most father's education was graduated from high school, and most mother's education was graduated from high school, the father's occupation mainly was self-employed, and the mother's job was not working (as a housewife).

Table 2: Distribution of the characteristics of the respondents' parents

Parental characteristics	f	%
Age (mean \pm SD) (years)		
Mother	30.05 \pm 6.18	
Father	34.55 \pm 6.91	
Mother education:		
Not schooling	12	3.7
Not completing the elementary school	4	1.2
Graduated from elementary school	30	9.3
Not finishing junior high school	6	1.9
Graduated from junior high school	43	13.3
Not finishing high school	15	4.6
Graduated from high school	140	43.2
Completed the university education	74	22.8
Father education:		
Not schooling	5	1.5
Not completing the elementary school	8	2.5
Graduated from elementary school	26	8.0
Not finishing junior high school	5	1.5
Graduated from junior high school	47	14.5
Not finishing high school	13	4.0
Graduated from high school	155	47.8
Completed the university education	65	20.8
Mother occupation:		
No work	221	68.2
Farmer	13	4.0
Labor	3	0.9
Entrepreneur	34	10.8
Civil servants	26	8.0
Traders	11	3.4
Private employees	15	4.6
Father occupation:		
No work	9	2.8
Farmer	65	20.1
Breeder	1	0.3
Labor	18	5.6
Entrepreneur	159	49.1
Civil servants	38	11.7
Traders	8	2.5
Private employees	26	8

Table 3 shows that most of the samples (84.8%) had a very good early breastfeeding practices status. About 93.2% of the samples were very good at exclusive breastfeeding. In comparison, 81.8% of the samples had a very good breastfeeding duration. The highest sample energy intake was in the fairly good category (36.7%). The most protein consumption was in the very good category (84.0%). Most samples had Vitamin A in the very good category (67.0%). The highest sample of Fe intake was in a low category (46%). The highest category of iodine intake was very low (82.1%). Nutritional status by bodyweight/age index

Table 3: Distribution of samples according to breastfeeding, consumption, and nutritional status

Indicators	n	%
Breastfeeding		
Early breastfeeding		
Good (breastfed immediately after birth)	273	84.3
Less (indirect breastfeeding was born)	51	15.7
Exclusive breastfeeding		
Good (6 months)	302	93.2
Less (<6 months)	22	6.8
Duration of breastfeeding		
Good (2 years)	265	81.8
Less (<2 years)	59	18.2
Energy Consumption		
Good (1300 sRDA=1400) kcal	119	36.7
Less (RDA<1300) kcal	205	63.3
Protein Consumption		
Good (20sRDA=25) g	272	84.0
Less (RDA <20) g	52	16.0
Vitamin A Intake		
Good (400sRDA=450) RE	217	67.0
Less (RDA<400) RE	107	33.0
Fe Intake		
Good (7sRDA=10) mg	149	46.0
Less (RDA<7) mg	175	54.0
Iodine Intake		
Good (90sRDA=120)mcg	58	17.9
Less (RDA<90) mcg	266	82.1
Zinc Intake		
Good (3sRDA=5) mg	172	53.1
Less (RDA<3) mg	152	46.9
Bodyweight / Age		
Good (-2sSD<2)	304	93.8
Less (<-2 SD)	20	6.2
Bodyheight / Age		
Good (-2sSD<2)	267	82.4
Less (SD<-2)	57	17.6
Bodyweight/ Bodyheight		
Good (-2sSD<2)	305	94.1
Less (SD<-2)	19	5.9
BMI / Age		
Good (-2sSD<1)	307	94.8
Less (SD<-2)	17	5.2

was 93.8% samples classified good categories, height/age index was 82.4% sample in good categories, index weight/height was 94.1% sample in the good category, and BMI/Age index 94.8% was a very good category.

Analysis of nutritional status indicators (NS1, NS2, NS3, and NS4) was said to be able to explain the nutritional status variable if it had a critical ratio (C.R.) > 1.96 or a $p < 0.05$. In other words, it can also be seen that the estimated loading value was > 0.5. Table 4 shows that of the four indicators, three indicators had an estimate loading value > 0.5, namely NS1, NS3, and NS4. It means that the three indicators were valid to explain the nutritional status variables used in model analysis. The breastfeeding indicator shows the three indicators, and all indicators had an estimate loading value of > 0.5, namely BF1, BF2, and BF3. It means that the three indicators were valid to explain breastfeeding variables to be used in model analysis. The consumption indicator shows the six indicators. All of them were valid to explain the consumption variable used in the model analysis (Table 4).

The measurement model is part of the SEM model, consisting of a dependent variable or several independent variables and several indicator variables from each of these variables. After the measurement model was proven valid, the process was continued by carrying out a structural model, analyzing the relationship between the dependent and independent variables and between indicator variables.

Table 4: Confirmatory factor analysis indicators of nutritional status, breastfeeding, and consumption

Indicators	Estimate	S.E.	C.R. ^{***}	p-value
NS1**	1.000			
NS2*	-0.579	0.458	-1.266	0.205
NS3**	4.800	1.385	3.466	***
NS4**	3.884	1.011	3.841	***
BF1**	1.000			
BF2**	1.171	0.279	4.202	***
BF3**	1.296	0.421	3.079	0.002
CF2	0.721	0.100	7.242	***
CF3	1.072	0.135	7.922	***
CF4	1.099	0.128	8.556	***
CF5	0.707	0.126	5.626	***
CF6	0.279	0.085	3.291	0.001
CF1	1.000			

*Not valid, **Valid, ***CR: Critical ratio

Figure 1 illustrates the relationship between the dependent and between independent and indicator variables. SEM model looked at the dependent variable nutritional status (NS1, NS3, and NS4) and the independent variable consumption (CF1, CF2, CF3, CF4, CF5, and CF6), breastfeeding (BF1, BF2, BF3).


**Figure 1: Respecification model of the relationship between dependent variables on nutritional status, independent variable of breastfeeding, and indicator variables**

Table 5 shows the result based on the two methods of the model's feasibility test, FGI (Goodness of Fit Index) ≥ 0.90 and AGFI (Adjusted Goodness of Fit Index) ≥ 0.80 . It shows that the goodness of fit index results of the model respecifications had met the required cut-off value, so the model was feasible (Table 5).

Table 5: Goodness of fit confirmatory index respecification model of nutritional status, consumption, and breastfeeding

Fit Index	Cut-off value	Model results	Note
FGI	≥ 0.90	0.974	Feasible
AGFI	≥ 0.80	0.659	Not Feasible

Table 6 shows with $p < 0.05$ or column estimate with a value > 0.5 shows the indicators BF1, BF2 and BF3. They could explain the independent variables of

Table 6: Results of loading regression weights relationship between nutritional status and breastfeeding with indicators

Dependent Variables, Independent and Indicators	Estimate	p-value
Nutrition Status <--- Breastfeeding	-0.005	0.830
Nutrition Status <--- Consumption	.013	0.438
NS1 <--- Nutrition Status	1.000	-
CF2 <--- Consumption	1.000	-
NS3 <--- Nutrition Status	4.818	0.000
NS2 <--- Nutrition Status	-6.466	0.000
NS4 <--- Nutrition Status	4.068	0.000
CF1 <--- Consumption	1.647	0.000
BF3 <--- Breastfeeding	1.195	0.000
BF2 <--- Breastfeeding	1.410	0.002
BF1 <--- Breastfeeding	1.000	-
CF6 <--- Consumption	.493	0.002
CF5 <--- Consumption	.883	0.000
CF4 <--- Consumption	1.686	0.000
CF3 <--- Consumption	1.882	0.000

breastfeeding. NS1, NS2, NS3, and NS4 indicators explained the dependent variable nutritional status. The indicators CF1, CF2, CF3, CF4, CF5, and CF6, explained the dependent variable consumption.

Table 7 shows breastfeeding and consumption factors explaining the weak relationship in nutritional status ($r = -0.015$) and ($r = 0.052$). If the value of $r = -0.015$ were squared, a value of 0.03% would be obtained. It means that the breastfeeding factor of 0.52% explained the nutritional status factor. The remaining 99.97% was explained by a unique factor, error factor (d1). For $r = 0.052$, it was 0.27%, means that the consumption factor was 0.27%. It explained the nutritional status factor, while the remaining 99.73 was explained by a unique factor, error factor (d1).

Table 7: Results of loading standardized regression weights: dependent, independent, and indicator variables

Dependent Variables and Indicators	Dependent and Independent Variables	Regression Coefficient (r)
Nutrition Status <--- Breastfeeding	Breastfeeding	-0.015
Nutrition Status <--- Consumption	Consumption	0.052
NS1 <--- Nutrition Status	Nutrition Status	0.203
CF2 <--- Consumption	Consumption	0.516
NS3 <--- Nutrition Status	Nutrition Status	0.997
NS2 <--- Nutrition Status	Nutrition Status	-0.982
NS4 <--- Nutrition Status	Nutrition Status	0.887
CF1 <--- Consumption	Consumption	0.647
BF3 <--- Breastfeeding	Breastfeeding	0.422
BF2 <--- Breastfeeding	Breastfeeding	0.764
BF1 <--- Breastfeeding	Breastfeeding	0.374
CF6 <--- Consumption	Consumption	0.244
CF5 <--- Consumption	Consumption	0.336
CF4 <--- Consumption	Consumption	0.680
CF3 <--- Consumption	Consumption	0.716

Discussion

1. The Relationship between Consumption Factors and Nutritional Status

Based on the confirmatory analysis factor, the consumption indicator ($r = 0.052$) or the consumption factor of 0.27% explained the nutritional status factor. The remaining 99.73 was defined by a unique factor, namely, error factor (d1). It causes the consumption factor to be further analyzed in the model structure analysis. In other words, the relationship between consumption factors can be analyzed directly with nutritional status.

Malnutrition is a severe health challenge factor in developing countries. Poor child feeding practices have been identified as a significant cause of stunting. Dietary diversity is one of the core indicators when assessing children's feeding practices and nutrition in obtaining nutrients that are essential for growth and development [16]. There were 102 foods important for children under five, namely (1) cereals, (2) spices condiments and beverages, (3) sweets, (4) oil and fats, (5) meat, (6) fruit, (7) vegetables, (8) milk and milk products, (9) tuber and roots, (10) legumes, nuts, and seeds, (11) eggs, (12) fish and other seafood [17]. Previous research had shown that most Indonesian children consume tuber vegetables and low consumption of fruit and dairy products [18].

Diversity in food consumption is a proxy for dietary quality and nutritional adequacy. Dietary variability is a problem for developing countries. Consumption of animal products is low, and there are few fruits and vegetables [19]. Furthermore, the staple plant foods consumed are poor in micronutrient content. The micronutrients they contain are also types that are difficult for the body to absorb [20]. The WHO recommends providing complementary foods for children starting at 6 months of age to meet the needs of calories, protein, and micronutrients needed for growth and development. Proper feeding during childhood is also crucial for preventing chronic degenerative diseases in the future [21].

This consumption factor is closely related to parenting by parents. According to UNICEF, parenting is defined as providing food, caring for children, guiding, and teaching children, which individuals and families do. The practice of giving food to children includes breastfeeding, quality supplementary food, food preparation and provision of nutritious food, child care, including caring for children when sick, immunization, giving supplements, bathing children, and so on [22]. The dominant factor that causes malnutrition is inappropriate behavior among the community in selecting and providing food to family members, especially children. Therefore, various activities must be carried out to provide proper nutrition (Feeding) and care (caring) to achieve good nutritional status. Feeding and caring through mothers' parenting to their children will positively and negatively affect their growth and development [23].

2. Relationship between factors of breastfeeding and nutritional status

The breastfeeding factor explains the weak relationship with nutritional status ($r = -0.015$), or the factor of breastfeeding of 0.03% explained the nutritional status factor. The remaining 99.97% was defined by a unique factor, namely, error factor (d1). It causes the consumption factor to be further analyzed in the model structure analysis. In other words, the relationship between breastfeeding factors could be analyzed directly with nutritional status [10].

Breastfeeding is a priority given to children [25]. Previous research shows that breastfeeding has a significant relationship with the incidence of stunting in children under five [24]. Breastfeeding is identified as an affordable maternal health behavior in protecting children against stunting. Breastfeeding can reduce the risk of 0.5 times the incidence of stunting. Breastfeeding for children is given at least until the age of 6 months to minimize the incidence of stunting in children [25].

Research that analyzed data from Riskesdas in 2013 found that exclusive breastfeeding was protective against stunting in children. However, the results were not significant, both for exclusive breastfeeding > 6 months and exclusive breastfeeding for 4- <6 months. The multivariable analysis results show that the variables associated with stunting in children aged 6-23 months in Indonesia are birth weight and household economic status. In the selected logistic regression equation model, children with low birth weight (<2,500 g) have a 1.77 times higher risk of becoming stunted compared to children born with normal weight (> 2,500 g) [26].

Weaning/stopping breastfeeding at an early age will impact the nutritional status of the child. Some of the factors that cause early weaning are the working mother's status, education level, and residence place [27]. Weaning has a significant role in determining a child's nutritional status. Poor weaning practices during infancy and toddlerhood can lead to malnutrition, contribute to cognitive and social growth and development disorders, poor school performance, and decreased productivity in the future [28].

Conclusion

The stunting model in children under five was associated with the analysis of factors of consumption and breastfeeding. Based on the confirmatory analysis, consumption and breastfeeding factors can be further analyzed with structural analysis models. The strength of the relationship between consumption and toddler stunting was relatively weak ($r = 0.052$). The strength of the relationship between breastfeeding and toddler stunting was relatively weak ($r = -0.015$). To prevent nutritional problems, the factors that need to be considered were consumption and breastfeeding. Other things that also need attention were the factors that cause the low level of consumption. It was particularly protein, and the things cause the mother's behavior to perform early breastfeeding practices and provide exclusive breastfeeding.

References

- Owino V, Ahmed T, Freemark M, Kelly P, Loy A, Manary M, *et al.* Environmental enteric dysfunction and growth failure/stunting in global child health. *Pediatrics*. 2016;138(6):e20160641. <https://doi.org/10.1542/peds.2016-0641> PMID:27940670
- Local Burden of Disease Child Growth Failure Collaborators. Mapping child growth failure across low-and middle-income countries. *Nature*. 2020;577(7789):231-4. <https://doi.org/10.1038/s41586-019-1878-8> PMID:31915393
- de Onis M, Branca F. Childhood stunting: A global perspective. *Matern Child Nutr*. 2016;12 Suppl 1:12-26. <https://doi.org/10.1111/mcn.12231> PMID:27187907
- Balitbangkes. Laporan Nasional Riskesdas 2013. Jakarta: Balitbangkes; 2014.
- Mahmudiono T, Sumarmi S, Rosenkranz RR. Household dietary diversity and child stunting in East Java, Indonesia. *Asia Pac J Clin Nutr*. 2017;26(2):317-25. PMID:28244712
- Alamu EO, Gondwe T, Eynia TE, Maziya-Dixon B. Assessment of dietary diversity of mothers and children of 6-24 months from eastern and Southern provinces of Zambia. *J Nutr Metab*. 2019;2019:1049820. <https://doi.org/10.1155/2019/1049820>
- Nachvak SM, Sadeghi O, Moradi S, Esmailzadeh A, Mostafai R. Food groups intake in relation to stunting among exceptional children. *BMC Pediatr*. 2020;20:1-8. <https://doi.org/10.1186/s12887-020-02291-7>
- Sirasa F, Mitchell L, Harris N. Dietary diversity and food intake of urban preschool children in North-Western Sri Lanka. *Matern Child Nutr*. 2020;16:1-17. <https://doi.org/10.1111/mcn.13006>
- Syeda B, Agho K, Wilson L, Maheshwari GK, Raza MQ. Relationship between breastfeeding duration and undernutrition conditions among children aged 0-3 years in Pakistan. *Int J Pediatr Adolesc Med*. 2020;8:10-7. <https://doi.org/10.1016/j.ijpam.2020.01.006> PMID:33718571
- Campos AP, Vilar-Compte M, Hawkins SS. Association between breastfeeding and child stunting in Mexico. *Ann Glob Health*. 2020;86(1):145. <https://doi.org/10.5334/aogh.2836> PMID:33262934
- Sari N, Manjorang MY, Zakiyah, Randell M. Exclusive breastfeeding history risk factor associated with stunting of children aged 12-23 months. *Kesmas*. 2021;16:28-32. <https://doi.org/10.21109/kesmas.v16i1.3291>
- Sirajuddin, Asbar R, Nursalim, Tamrin A. Breastfeeding practices can potential to prevent stunting for poor family. *Enfem Clin*. 2020;30 Suppl 4:13-7. <https://doi.org/10.1016/j.enfcl.2020.02.007> PMID:32545030
- Kemenkes RI. Laporan Nasional Riset Kesehatan Dasar 2018. Jakarta: Kementerian Kesehatan Republik Indonesia; 2019.
- TNP2K. 100 Kabupaten/Kota Prioritas Untuk Intervensi Anak Kerdil (Stunting). Jakarta: Tim Nasional Percepatan Penanggulangan Kemiskinan (TNP2K); 2017.
- Kemenkes RI. Angka Kecukupan Gizi yang Dianjurkan Bangsa Indonesia. Jakarta: Direktorat Bina Gizi dan Kesehatan Ibu dan Anak; 2014.
- Khamis AG, Mwanri AW, Ntwenya JE, Kreppel K. The influence of dietary diversity on the nutritional status of children between

- 6 and 23 months of age in Tanzania. *BMC Pediatr.* 2019;19:1-9. <https://doi.org/10.21203/rs.2.16657/v3>
17. Modjadji P, Molokwane D, Ukegbu PO. Dietary diversity and nutritional status of preschool children in North West Province, South Africa: A cross sectional study. *Children (Basel).* 2020;7(10):174. <https://doi.org/10.3390/children7100174> PMID:33050271
18. Kunto YS, Bras H. Ethnic group differences in dietary diversity of school-aged children in Indonesia: The roles of gender and household SES. *Food Nutr Bull.* 2019;40(2):182-201. <https://doi.org/10.1177/0379572119842993> PMID:31046454
19. Ahmad I, Khalique N, Khalil S, Urfi U, Maroof M. Dietary diversity and stunting among infants and young children: A cross-sectional study in Aligarh. *Indian J Community Med.* 2018;43(1):34-6. https://doi.org/10.4103/ijcm.ijcm_382_16 PMID:29531436
20. Berrazaga I, Micard V, Gueugneau M, Walrand S. The role of the anabolic properties of plant-versus animal-based protein sources in supporting muscle mass maintenance: A critical review. *Nutrients.* 2019;11(8):1825. <https://doi.org/10.3390/nu11081825> PMID:31394788
21. World Health Organization. WHA Global Nutrition Targets 2025: Stunting Policy Brief. Geneva: World Health Organization; 2014. Available from: https://www.who.int/nutrition/topics/globaltargets_stunting_policybrief.pdf. [Last accessed on 2021 Mar 16].
22. Tessema M, Gunaratna NS, Brouwer ID, Donato K, Cohen JL, McConnell M, et al. Associations among high-quality protein and energy intake, serum transthyretin, serum amino acids and linear growth of children in Ethiopia. *Nutrients.* 2018;10(11):1776. <https://doi.org/10.3390/nu10111776> PMID:30453477
23. Uwiringiyimana V, Ocké MC, Amer S, Veldkamp A. Predictors of stunting with particular focus on complementary feeding practices: A cross-sectional study in the northern province of Rwanda. *Nutrition.* 2019;60:11-8. <https://doi.org/10.1016/j.nut.2018.07.016> PMID:30508763
24. Barir B, Murti B, Pamungkasari EP. The Associations between exclusive breastfeeding, complementary feeding, and the risk of stunting in children under five years of age: A path analysis evidence from Jombang East Java. *J Matern Child Health.* 2019;4:486-98. <https://doi.org/10.26911/thejmch.2019.04.06.09>
25. Anindya IG, Salimo H, Dewi YL. Relationship between exclusive breastfeeding and stunting in children aged 6 months old. In: The 6th International Conference on Public Health; 2019. p. 213. <https://doi.org/10.26911/the6thicph.03.71>
26. Paramashanti BA, Hadi H. Exclusive breastfeeding practice was not related with stunting in young children 6–23 months in Indonesia. *J Gizi Diet Indones.* 2015;3:162-74. [https://doi.org/10.21927/ijnd.2015.3\(3\).162-174](https://doi.org/10.21927/ijnd.2015.3(3).162-174)
27. Pambudi J, Christijan R. Early weaning practice in correlation to social-economic and residential geography. *Penelit Gizi Makanan.* 2017;40:87-94. <https://doi.org/10.22435/pgm.v40i2.7666.87-94>
28. Vyas S, Kandpal SD, Semwal J, Chauhan SN. Trends in weaning practices among infants and toddlers in a hilly terrain of a newly formed state of India. *Int J Prev Med.* 2014;5(6):741-8. PMID:25013694

The Influence of Factors of Breastfeeding and Food Consumption on Stunting Incidence among Preschool Children in Keerom District, Papua Province, Indonesia

5
ORIGINALITY REPORT

19%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|---------------|
| 1 | www.scribd.com
Internet | 75 words — 2% |
| 2 | archives.palarch.nl
Internet | 53 words — 1% |
| 3 | Mona Hamdy, Iman Hassan Draz, Inas Talaat El Sayed, Azza Ali Fahmy Ayyad, Marwa Rashad Salemd. "Assessment of Quality of Life among Beta-Thalassemia Major Patients Attending the Hematology Outpatient Clinics at Cairo University Hospital", Open Access Macedonian Journal of Medical Sciences, 2021
Crossref | 51 words — 1% |
| 4 | www.science.gov
Internet | 49 words — 1% |
| 5 | ijcm.org.in
Internet | 47 words — 1% |
| 6 | eprints2.ipdn.ac.id
Internet | 38 words — 1% |
| 7 | Diane S. Kaplan, Xiaoru Liu, Howard B. Kaplan. "Explaining Dropout Behavior: A Longitudinal Study | 36 words — 1% |

of Direct and Indirect Psychosocial Effects of Seventh and Eighth Grade Experiences", Educational Research and Evaluation, 1995

Crossref

-
- 8 nveo.org 32 words — 1 %
Internet
-
- 9 Anwar Mallongi, Denny Adelyta Tofani Novitasari, Noer Bahry Noor, Fatmawati, Muh. Saleh, Sehalyana, Apollo, Rosmala Nur. "Health risk analysis of phenol and arsenic exposure among kokoda communities village Sorong, West Papua", Enfermería Clínica, 2020 28 words — 1 %
Crossref
-
- 10 Perpetua Modjadji, Dineo Molokwane, Patricia Ogechi Ukegbu. "Dietary Diversity and Nutritional Status of Preschool Children in North West Province, South Africa: A Cross Sectional Study", Children, 2020 27 words — 1 %
Crossref
-
- 11 repository.uki.ac.id 27 words — 1 %
Internet
-
- 12 www.nutriweb.org.my 27 words — 1 %
Internet
-
- 13 repository.unism.ac.id 19 words — < 1 %
Internet
-
- 14 www.dovepress.com 19 words — < 1 %
Internet
-
- 15 zombiedoc.com 19 words — < 1 %
Internet

16 Ahmed Gharib Khamis, Akwilina Wendelin Mwanri, Julius Edward Ntwenya, Katharina Kreppel. "The influence of dietary diversity on the nutritional status of children between 6 and 23 months of age in Tanzania", BMC Pediatrics, 2019

Crossref

18 words — < 1%

17 Meitria Syahadatina Noor, Ayu Riana Sari, R. Akbar Agustriyanto, Rezeki Norwinardi et al. "Role of Cadre in Improving Knowledge and Attitude of Chronic Energy Deficiency on Teenagers in Mali-Mali Village, Banjar Regency, South Kalimantan, Indonesia", Open Access Macedonian Journal of Medical Sciences, 2021

Crossref

15 words — < 1%

18 Ni Wayan Dewi Tarini, Moses Glorino Rumambo Pandin. "PRACTICE OF EARLY BREASTFEEDING INITIATION ON PRIMIGRAVIDA WITH A CULTURAL PERSPECTIVE: A SYSTEMATIC REVIEW", Cold Spring Harbor Laboratory, 2022

Crossref Posted Content

15 words — < 1%

19 garuda.ristekbrin.go.id

Internet

15 words — < 1%

20 ndl.ethernet.edu.et

Internet

15 words — < 1%

21 Hsiang-Ann Liao. "Examining the Role of Collaborative Learning in a Public Speaking Course", College Teaching, 2014

Crossref

14 words — < 1%

22 Sukmawati Sukmawati, Yanti Hermayanti, Eddy Fadlyana, Henny Suzana Mediani. "Stunting Prevention with Education and Nutrition in Pregnant Women: A Review of Literature", Open Access Macedonian Journal of

14 words — < 1%

Medical Sciences, 2021

Crossref

-
- 23 bmcmededuc.biomedcentral.com¹⁰
Internet 14 words — < 1%
-
- 24 repository.ub.ac.id
Internet 14 words — < 1%
-
- 25 Atika Dranesia, Dessie Wanda, Happy Hayati.
"Pressure to eat is the most determinant factor
of stunting in children under 5 years of age in Kerinci region,
Indonesia", Enfermería Clínica, 2019
Crossref 12 words — < 1%
-
- 26 jgp.poltekkes-mataram.ac.id
Internet 12 words — < 1%
-
- 27 Ahmed Yamany Ali, Abeer Attia Abdelkhalik,
Shereen Esmat, Walaa Alsharany Abuelhamd,
Hend Abdullah Elshemy, Mervat Essam Behiry. "Awareness and
Attitude toward Infection Control among Health Care Workers
during Coronavirus Disease-19 Pandemic, Egypt", Open Access
Macedonian Journal of Medical Sciences, 2021
Crossref 11 words — < 1%
-
- 28 faculty.uobasrah.edu.iq
Internet 11 words — < 1%
-
- 29 Aldina Ayunda Insani, Firdawati, Wella Dwi Anjani.
"RELATIONSHIP BETWEEN EXCLUSIVE
BREASTFEEDING WITH STUNTING EVENTS IN AGE 12-35
MONTHS IN AIR DINGIN HEALTH CENTER PADANG 2018",
[Walter de Gruyter GmbH](https://www.walterdegruyter.com)⁶, 2020
Crossref 10 words — < 1%
-
- 30 ejournal.almaata.ac.id⁶
Internet 10 words — < 1%

10 words — < 1%

31

www.atlantis-press.com
Internet

10 words — < 1%

32

¹³
⁴ Aisyiyah Alviana Agustin, Muhammad Ali Shodikin.
"Risk Factors for Stunting at Balung Health Center,
Jember Regency, Indonesia", Jurnal Aisyah : Jurnal Ilmu
Kesehatan, 2021

Crossref

9 words — < 1%

33

Gina Kennedy, Guy Nantel, Inge D Brouwer, Frans J
Kok. "Does living in an urban environment confer
advantages for childhood nutritional status? Analysis of
disparities in nutritional status by wealth and residence in
Angola, Central African Republic and Senegal", Public Health
Nutrition, 2007

Crossref

9 words — < 1%

34

Marion L. Roche. "Infant and young child feeding in
the Peruvian Amazon: the need to promote
exclusive breastfeeding and nutrient-dense traditional
complementary foods", Maternal and Child Nutrition, 02/2010

Crossref

9 words — < 1%

35

William Bellotti, Esta Lestari, Karen Fukofuka. "A
Food Systems Perspective on Food and Nutrition
Security in Australia, Indonesia, and Vanuatu", Elsevier BV, 2018

Crossref

9 words — < 1%

36

biointerfaceresearch.com
Internet

9 words — < 1%

37

care.diabetesjournals.org
Internet

³
9 words — < 1%

38 journal.poltekkes-mks.ac.id 9 words — < 1%
Internet

39 repo.unand.ac.id 9 words — < 1%
Internet

40 s-space.snu.ac.kr 9 words — < 1%
Internet

41 ¹²[Iqlima Dwi Kurnia, Ilya Krisnana, Tyas Dwi](#) 8 words — < 1%
⁸[Rahmadhani. "Perceived barriers of nutritional feeding related to severe stunting", Enfermería Clínica, 2021](#)
Crossref

42 discovery.researcher.life 8 words — < 1%
Internet

43 Hans Antlöv, Anna Wetterberg, Leni Dharmawan. 7 words — < 1%
"Village Governance, Community Life, and the 2014
⁹[Village Law in Indonesia", Bulletin of Indonesian Economic Studies, 2016](#)
Crossref

44 Mikawati ., Evi Lusiana, Hasriany .. "The 7 words — < 1%
Relationship between Exclusive Breastfeeding (ASI)
and Mother Heightwith Incident Rates Stunting among Child
Age 2-5 Years In Barombong Public Health Center, Gowa,
Sulawesi Selatan", KnE Life Sciences, 2019
Crossref

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF

Breastfeeding and Consumption Food

ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

8%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

1	www.x-mol.com Internet Source	2%
2	Scott E. Lewis. "Retention and Reform: An Evaluation of Peer-Led Team Learning", Journal of Chemical Education, 2011 Publication	2%
3	Submitted to Soongsil University Student Paper	2%
4	aisyah.journalpress.id Internet Source	2%
5	Submitted to Universitas Jenderal Achmad Yani Student Paper	1%
6	digilib.iain-palangkaraya.ac.id Internet Source	1%
7	Submitted to Ateneo de Manila University Student Paper	1%
8	Iqlima Dwi Kurnia, Ilya Krisnana, Tyas Dwi Rahmadhani. "Perceived barriers of	1%

nutritional feeding related to severe stunting", Enfermería Clínica, 2021

Publication

9	ilo.ch Internet Source	1 %
10	Submitted to Bogazici University Student Paper	1 %
11	Submitted to Universitas Bengkulu Student Paper	1 %
12	www.elsevier.es Internet Source	1 %
13	Aisiyiah Alviana Agustin, Muhammad Ali Shodikin. "Risk Factors for Stunting at Balung Health Center, Jember Regency, Indonesia", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2021 Publication	1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

Breastfeeding and Consumption Food

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14