

# Malaria among young children in Twano-Papua: Evidence of determining factors in endemic areas

Ruth Yogi<sup>1\*</sup>, Edison Kabak<sup>2</sup>

<sup>1</sup>Midwifery Study Program, Health Polytechnic Ministry of Health of Jayapura

<sup>2</sup>Nursing Study Program, Health Polytechnic Ministry of Health of Jayapura

## ARTICLE INFO

### Article history:

Received Mar 18, 2023

Revised Mar 28, 2023

Accepted Jun 21, 2023

### Keywords:

Children under 5 years  
malaria  
Papua  
Twano

## ABSTRACT

Malaria that occurs in children under 5 years has a high risk. Therefore, malaria control is made one of the main priorities in the health program by the Government. It is not yet known what the determining factors related to the incidence of young children in the Twano of Jayapura area. The objective is to identify the factors determinant associated with the incidence of malaria in young children (aged 0-5 years) in the working area of Twano of Jayapura. This study used a quantitative research with cross-sectional study. It was held in January 2021 with a sample size of 55 participated out of 61 people total samples. The inclusion criteria were mothers who had children aged 0-5 years. We used non-probability sampling technique. Data were obtained through questionnaires arranged according to the Likert Scale. Data were analyzed using SPSS version 25 with Chi-Square statistical test. The results showed three determinant factors associated with the incidence of malaria in children aged 0-5 years in Twano, Jayapura namely mothers' knowledge (p-value of 0.003), attitude (p-value of 0.005), and environment (p-value of 0.000). While the factors that were not associated with the incidence of malaria in children aged 0-5 years were education mothers (p-value of 0.318) and socioeconomic (p-value of 0.369). We found three important issues that influence the occurrence of malaria cases as determining factor in young children in Twano, Jayapura, namely knowledge, attitudes, and the environment. public health field practices based on the 2019 National Guidelines for HIV/AIDS Management.

This is an open access article under the [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) license.



## Corresponding Author:

Ruth Yogi,  
Midwifery,  
Poltekkes Kemenkes Jayapura,  
Padang Bulan, RW.2, Hedam, Kec. Heram, Jayapura City, Papua 99351  
Email: [ruthyogi8@gmail.com](mailto:ruthyogi8@gmail.com)

## INTRODUCTION

The World Health Organization (WHO) in 2020 estimated 241 million cases of malaria with an annual mortality rate of 627,000 deaths (World Health Organization., 2021). The incidence of malaria decreased from 81 in 2000 to 59 in 2015 and 56 in 2019, before increasing to 59 in 2020 (Konaté et al., 2020). The Ministry of Health recorded that the total malaria cases in Indonesia in

2020 were 254,055 (Ipa, Laksono, et al., 2020; Ipa, Widawati, et al., 2020). The percentage of malaria suspects with microscopic laboratory confirmation and the Rapid Diagnostic Test (RDT) in 2020 is 97% with a total of 1,823,104 examinations from 1,877,769 suspects with a PR (positivity rate) of 14% (Aprilia et al., 2020). About one-third of the incidence of malaria occurs in young children, namely children aged 0-5 years (Patriani et al., 2019). The incidence of malaria in children aged 0-5 years will result in impaired growth and stunted pubertal development (Ouédraogo et al., 2018). Based on the achievement of endemicity per province in 2020, 3 provinces have achieved 100% malaria elimination, including DKI Jakarta, East Java, and Bali (Hasyim et al., 2019). Meanwhile, the provinces with areas that have not yet achieved malaria elimination are Maluku, Papua, and West Papua (Budiarti et al., 2020; Dhewantara et al., 2019).

As part of its global commitment to eliminating malaria, the Indonesian government has issued a Minister of Health Decree to strengthen malaria control efforts from the central level to Puskesmas (Istiana et al., 2021). Elimination refers to efforts to stop the transmission of local (indigenous) malaria in a certain geographic area (Guntur, Kingsley, & Amirul Islam, 2021; Guntur, Kingsley, & Islam, 2021; Oyibo et al., 2021). Therefore, vigilance is still needed to prevent re-infection in areas that are considered malaria-free, including malaria in children under the age of 5 years. The biggest obstacles to eliminating malaria, according to the 2017 World Malaria Report, are the emergence of parasite resistance to antimalarial drugs, mosquito resistance to insecticides, and inadequate health system performance (Morakinyo et al., 2018; Tizifa et al., 2018). The results of further analysis of the 2013 Basic Health Research with household analysis units in 6 malaria-endemic provinces in Indonesia showed that most of the anti-malarial drugs were obtained by households from pharmacies and formal health services (Hasyim et al., 2019; Ipa, Widawati, et al., 2020). However, there are not a few studies that discuss other determining factors such as the role of mothers, education, nutrition, and the household environment. The government targets Indonesia to be free of malaria by 2030 (Rahmasari et al., 2021).

One of the efforts to achieve this target is determined by not only the effectiveness of treatment but also the integral role of various sectors (Bandzuh et al., 2022; Guntur, Kingsley, & Amirul Islam, 2021). Malaria in Indonesia remains difficult to eradicate even though the cause is known, there is control, and the cure has been given to patients because the complexity of an archipelagic country with diverse access to health services contributes to the progress of achieving malaria elimination (Istiana et al., 2021). In Papua, there are four districts/cities experiencing malaria endemic, namely Jayapura City, Jayapura Regency, Merauke, and Timika (Debora et al., 2018; Manangsang et al., 2021). There are many obstacles to achieving Malaria-Free Indonesia 2030, including people who think that malaria is a common disease, so they don't pay much attention (Oyibo et al., 2021; Rizkyansah & Rahayu, 2021). Research on the role of mothers in the occurrence of malaria in children under the age of 5 years in endemic areas has not been widely studied in Papua (Debora et al., 2018; Patriani et al., 2019). The gap in this research is what distinguishes it from previous studies which are expected to contribute new ideas for the management of malaria in Papua in particular and endemic areas in general.

Many studies about malaria focus on control efforts include self-protection against mosquito bites by using insect repellants, protective clothing, mosquito coils, mosquito nets, and insect sprays, and maintaining environmental cleanliness including draining, burying, and closing water storage containers and plants that can repel mosquitoes in the area and home environment (Hasyim et al., 2018; Morakinyo et al., 2018). That is what distinguishes this study from the fact that it focuses more on the determinants of malaria in children in endemic areas. Some of the determining factors are mothers with high knowledge make provision to think before taking action to prevent the incidence of malaria in their children (Ouédraogo et al., 2018). The action or behavior of a clean and healthy life carried out at home or in the home environment is also considered to play a role as a determining factor (Aberese-Ako et al., 2019). Malaria cases in Papua are the highest Annual Parasite Incidence (API), which are 123.00 per 1.000 population

with a positive incidence of malaria in as many as 137,265 (31,74%) (Sroyer et al., 2022). The incidence of infants and toddlers suffering from malaria in Papua Province is 13,832 (10.05%) (Debora et al., 2018). The number of malaria sufferers in 2018 was 20,119 cases and an increase in 2019 of 22,045 cases and 2,204 (10%) malaria cases occurred in infants and toddlers (Dinas Kesehatan Prov Papua 2020).

This quantitative study with a cross-sectional design approach tried to identify the determinants of malaria that occur in young children in Twano, in Jayapura City. The number of malaria cases in 2018 was 58 cases and in 2019 there were 108 cases (Kemenkes RI, 2019). It is not yet known whether the increase in malaria cases is related to the role of knowledge, attitudes, and the environment pf mothers. Therefore, this study aims to identify the determinant factors associated with the incidence of malaria in children aged 0-5 years in the region.

## RESEARCH METHODS

This quantitative research used a cross-sectional approach with a descriptive design. The research location is in Twano, one of the endemic areas in Jayapura, eastern Indonesia. The study population was mothers with children under the age of 5 years (young children). The number of samples was 61 people. Those who participated in the study were 55 people (90.2%) who according to WHO had met the requirements because they were more than 85% (Guntur, Kingsley, & Islam, 2021; Mahkota et al., 2020). We used a questionnaire validated with 4 parts of the list of questions. The first part is about demographic data which contains information on gender, age, education level, occupation, family size, and household socioeconomic. The second part is about general knowledge of mothers about malaria including symptoms, causes, and prevention. Our third section collects information on when, and where to take treatment for malaria, and the protection of children under 5 years of age. The fourth section is about the environment, the main parts of the house, access to drinking water, the closest health facility, and the distance to the nearest health facility. The research variables included the independent (education, knowledge, attitude, socioeconomic and environmental) and the dependent variable (the incidence of malaria in young children). Descriptive analysis was conducted to show the distribution of the characteristics of different respondents. The data were analyzed using SPSS and to evaluate the relationship between sociodemographic and environmental characteristics, we applied the chi-square test. We complied with the Declaration of Helsinki to ensure the protection of the rights, integrity, and confidentiality of respondents. All respondents signed the consent form before the interview.

## RESULTS AND DISCUSSION

In this section, we will describe the demographic data, education and knowledge, attitude, social economy, and environmental conditions, as well as discussing the day-to-day results of data analysis.

### Demographic Data

**Table 1.** Mothers' age and Employment Status

Age	N	%	Employment	N	%
18-25	20	36.4	Employed	32	58.2
26-35	28	50.9	Unemployed	23	41.8
>35	7	12.7			
Total	55	100		55	100

Table 4.1 shows that of the 55 mothers, most of them were 26-35 years old, as many as 28 people (50.9%), and a few aged <20 years as many as 3 people (5.5%). Their employment status is mostly not working as many as 32 people (58.2%).

### Education and knowledge

**Table 2.** Mothers' Education Status and Knowledge on Malaria

Education	N	%	Knowledge on Malaria	N	%
Lower	19	34.5	Less	21	38.2
High	36	64.5	good	34	61.8
Total	55	100		55	100

Table 2 shows that of the 55 mothers, most of them have higher education as many as 36 people (65.5%) and 34 people (61.8%) have good knowledge of malaria.

### Attitude, social economy, and environmental conditions

**Table 3.** Mothers' Attitude towards Malaria, social economic status, and environmental condition

Attitude	N	%	Social-economy status	N	%	Environment condition	N	%
negative	12	21.8	Low	29	52.7	Bad	20	36.4
High	43	78.2	Medium	26	47.3	good	35	63.6
Total	55	100		55	100	Total	55	100

Table 3 shows that most of the mothers have a positive attitude (43 people or 78.2%) with poor socioeconomic status (29 people or 52.7%). Good environmental conditions for 35 people (63.6%) and poor environmental conditions for as many as 20 people (36.4%).

### Bivariate Analysis

**Table 4.** Correlation between mothers' education and Malaria among young children

Knowledge	among young children				Total		<i>p-value</i>
	Malaria cases among children				N	%	
	Positive	Negative					
	n	%	n	%			
Less	12	57.1	9	42.9	21	100	0.003
good	5	14.7	29	85.3	34	100	
Total	17	30.9	38	69.1	55	100	

Table 4 shows the results of the chi-square statistical test at a significance value of 95% ( $\alpha=0.05$ ) obtained a *p*-value of 0.318 or  $p>(0.05)$ . Thus, there is no significant relationship between education and the incidence of malaria in young children in Twano, Jayapura City.

**Table 5.** Correlation between Mothers' Knowledge and Malaria Among Young Children

Knowledge	Malaria cases among young children				Total		<i>p-value</i>
	Positive		Negative		N	%	
	n	%	n	%			
Less	12	57.1	9	42.9	21	100	0.003
Good	5	14.7	29	85.3	34	100	
Total	17	30.9	38	69.1	55	100	

Table 5 shows the results of the chi-square statistical test at a significance value of 95% ( $\alpha=$

0.05) obtained a p-value of 0.003 or  $p < (0.05)$ . Thus, there is a significant relationship between knowledge and the incidence of malaria among young children in Twano, Jayapura City.

**Table 6.** Correlation between mothers' attitude and Malaria among young children

Table 6: Correlation between mothers' attitude and malaria among young children							
Attitude	Malaria cases among young children				Total		<i>p-value</i>
	Positive		negative		N	%	
	n	%	n	%			
negative	8	66.7	4	33.3	12	100	0.005
Positive	9	20.7	34	79.1	43	100	
Total	17	30.9	38	69.1	55	100	

Table 6 show the results of the chi-square statistical test at a significance value of 95% ( $\alpha = 0.05$ ) obtained a p-value of 0.005 or  $p < (0.05)$ . Thus, there is a significant relationship between attitudes and the incidence of malaria in young children in Twano, Jayapura City.

**Table 7.** Correlation between social economic status and Malaria among young children

Social economy status	among young children				Total		<i>p-value</i>
	Malaria cases among young children				N	%	
	Positive	Negative					
	n	%	n	%			
Low	11	37.9	18	62.1	29	100	0.369
Medium	6	23.1	20	76.9	26	100	
Total	17	30.9	38	69.1	55	100	

Table 7 show the results of the chi-square statistical test at a significance value of 95% ( $\alpha = 0.05$ ) obtained with a p-value of 0.369 or  $p > (0.05)$ , thus there is no significant socio-economic relationship with the incidence of malaria in young children in Twano, Jayapura City.

**Table 8.** Correlation between environmental conditions and Malaria among young children

Environment conditions	Malaria cases among young children				Total		<i>p-value</i>
	Positive		Negative		N	%	
	n	%	n	%			
Bad	13	65	7	35	20	100	0.000
good	4	11.4	31	88.6	35	100	
Total	17	30.9	38	69.1	55	100	

Table 8 show the results of the chi-square statistical test at a significance value of 95% ( $\alpha = 0.05$ ) obtained with a p-value of 0.000 or  $p < (0.05)$ , thus there is a significant environmental relationship with the incidence of malaria in children aged 0-5 years at the Twano, Jayapura City.

## Discussion

We found three important issues that influence the occurrence of malaria cases as a determinant factor in young children in Twano, Jayapura, namely knowledge, attitude, and environment.

## Knowledge

The results of the study were 12 people (57.1%). Our analysis obtained a p-value of 0.003 or  $p < (0.05)$ , thus there is a significant correlation between knowledge and the incidence of malaria in children aged 0-5 years at Twano of Jayapura City. Knowledge is something that exists or is considered to exist, something that is the result of conforming the subject with the object, the result of human nature wanting to know, the result of the compatibility between induction and

deduction, as a picture of external objects that are present in the human mind and something present and materialized (Sembiring & Lubis, 2019). Mothers' knowledge is very influential on the actions taken in preventing malaria in their children (Olubiyi, Folami, 2018). Knowledge can include how to prevent malaria in their children, for example at the time of biting malaria and how malaria transmission occurs in their children. Lack of knowledge of mothers is found especially in mothers who have low education and lack of experience about malaria which has an impact on malaria prevention actions taken by mothers to their children (Yasuoka et al. 2018). Lack of knowledge has an impact on maternal actions in preventing malaria.

### **Attitude**

The results of our study on mothers who have a negative attitude with a positive incidence of malaria in young children we found 8 people (66.7%). Research results obtained a p-value of 0.005 or  $p < (0.05)$ , thus there is a significant correlation between attitudes and the incidence of malaria in young children in Twano, Jayapura City. This shows that a negative attitude causes mothers to be less able to prevent malaria because malaria is a disease that often occurs in the community (Oluwasogo; Henry; Abdurashed; Olawuni; and Olabisi, 2016). Attitude is a person's closed response to a stimulus or object, both internal and external, so that its manifestation cannot be directly seen, but can only be interpreted beforehand from the closed behavior (Orhan & Serin, 2019). Attitudes, in reality, indicate the suitability of the response to a particular stimulus (Herman, 2020). A lot of research revealed attitudes that had less impact on the mother's lack of action in preventing malaria because she consider that malaria is an endemic disease that often occurred in her area (Saha et al., 2019). The level of public awareness of the dangers of malaria can affect the public's willingness to take preventive measures to overcome the possibility of contracting malaria (Padonou et al., 2018). Public awareness can be seen in the preventive measures taken such as the habit of being out of the house until late at night, carrying out environmental health activities, and using mosquito nets (Rahmasari et al., 2021).

Mothers who have a negative attitude toward the negative incidence of malaria in young children in this study were 4 people (33.3%). This is because even though the mother has a poor attitude, in prevention, they practices good prevention methods for their child (Patriani et al., 2019). For example, in the afternoon, the children are bathed and prohibited from going out to play. In addition, mothers who have good socio-economic conditions can fulfill good home sanitation facilities in preventing mosquitoes from entering the house. However attitudes cannot be automatically manifested in everyday actions. The existence of supporting factors such as facilities and support is needed to turn attitudes into real actions.

On the other hand, there were 34 mothers (79.1%) who had a positive attitude with a negative incidence of malaria. This means that the level of a good mother's attitude tends to be good and eventually behaves well in preventing the incidence of malaria in her child. A good mother's attitude are shown by keeping her home environment clean, installing rough wires, and preventing mosquito bites on her child when going out at night by wearing long sleeves. A good attitude will affect the incidence of malaria because it will have an impact on the health of the family if they do not do good prevention (Roosihermatie et al., 2017). Other studies have shown that an individual's health status is also influenced by behavioral factors, namely the attitude of a person who has a major impact on a person's health status (Afoakwah et al., 2018; Yasuoka et al., 2018).

### **Environment**

The results of this study indicate mothers who have a poor environment with a positive incidence of malaria in children aged 0-5 years are 13 people (65%). The analysis obtained a p-value of 0.000 or  $p < (0.05)$ , thus there is a significant environmental relationship with the incidence of malaria in young children in Twano, Jayapura. Environmental factors that are risk factors for malaria transmission are the presence of stagnant water which can be a breeding habitat for

Anopheles mosquitoes, the presence of dense bushes around people's residences and close to the Anopheles breeding habitat, house ventilation is not covered with gauze, does not have a house ceiling, and the walls of the house are not tight (Wanzira et al., 2017). Environmental conditions affect the breeding of mosquitoes so that children are susceptible to the incidence of malaria (Aberese-Ako et al., 2019). An unfavorable residential environment poses an increased risk of malaria transmission because there are still many Anopheles mosquito breeding sites (Abossie et al., 2020; Tizifa et al., 2018). This is due to the environment of the mother who lives around the swamp area in the Hamadi and Entrop areas, Jayapura which is a breeding ground for mosquitoes and mothers who are not paying attention to their home environment such as installing wire screens, no damaged ceilings or ceilings, lack of lighting causes mosquitoes can easily enter the house (Roosihermatie et al., 2017). Residents who are not healthy have a higher chance of malaria incidence compared to healthy settlements (Hasyim et al., 2018). Other aspects cannot be fulfilled by the mother's family, such as having a good ceiling, but mothers can take precautions by diligently cleaning the inside and outside of the house and draining and closing the water reservoir so that mosquitoes cannot breed in the house. Healthy environmental conditions can be realized by the existence of healthy community behavior so that it can reduce the presence of disease vectors in the house (Bandzuh et al., 2022). Healthy environmental conditions can be realized by the existence of healthy community behavior.

#### **Study Limitation**

The limitation of this study is that the results cannot be used for cause analysis, considering that research and assessment in cause analysis require a clear time sequence between exposure and disease, i.e. exposure precedes disease. This research also only took place at a certain point in time, was not followed up, and did not involve variable manipulation. Many malaria studies have been carried out in Papua, but no malaria cases have been reported among young children there as a malaria-prone area of Twano. This gap is what distinguishes it from previous studies that contributed novelty to research on malaria in Papua (Dhewantara et al., 2019; Inriyanti et al., 2021).

## **CONCLUSION**

The purpose of this study was to identify the determinant factors associated with the incidence of malaria in children aged 0-5 years in the Twano area, Jayapura City. After identifying cases in the field. We found three important issues that influence the occurrence of malaria cases as a determinant factor in young children in Twano, Jayapura, namely knowledge, attitude, and environment. The solution requires a structured approach, not only intensive individual health education but also a multi-sector approach, from the local government, health workers, social services, and sanitation. Everything is mutually supportive and integrated. Therefore, in the future, we further research on the management of determinant factors of malaria in endemic areas involving multi-professionals particularly in Twano in particular and Papua in general is highly recommended.

## **ACKNOWLEDGEMENT**

The authors acknowledged the Poly Technic of Health of the Ministry of Health of Jayapura for supporting the study, and our gratitude to all participants who were willing to participate in the study.

## **References**

- Aberese-Ako, M., Magnussen, P., Ampofo, G. D., & Tagbor, H. (2019). Health system, socio-cultural, economic, environmental and individual factors influencing bed net use in the prevention of malaria in

- pregnancy in two Ghanaian regions. *Malaria Journal*, 18(1), 1-13. <https://doi.org/10.1186/s12936-019-2994-5>
- Abossie, A., Yohanes, T., Nedu, A., Tafesse, W., & Damitie, M. (2020). Prevalence of malaria and associated risk factors among febrile children under five years: A cross-sectional study in arba minch zuria district, south Ethiopia. *Infection and Drug Resistance*, 13, 363-372. <https://doi.org/10.2147/IDR.S223873>
- Afoakwah, C., Deng, X., & Onur, I. (2018). Malaria infection among children under-five: The use of large-scale interventions in Ghana. *BMC Public Health*, 18(1), 1-13. <https://doi.org/10.1186/s12889-018-5428-3>
- Aprilia, F., Faraid, S., Handayani, I., & Esa, T. (2020). Profile of Rapid Molecular Test of Tuberculosis Using Xpert MTB/RIF Profile of Rapid Molecular Test Faraid-, et al. 26(2), 223-228. [www.indonesianjournalofclinicalpathology.org](http://www.indonesianjournalofclinicalpathology.org)
- Bandzuh, J. T., Ernst, K. C., Gunn, J. K. L., Pandarangga, S., Rambu, L., Yowi, K., Hobgen, S., Cavanaugh, K. R., Kalaway, R. Y., Rambu, N., Kalunga, J., Felipus Killa, M., Ara, U. H., Uejio, C. K., & Hayden Id, M. H. (2022). Knowledge, attitudes, and practices of Anopheles mosquito control through insecticide treated nets and community-based health programs to prevent malaria in East Sumba Island, Indonesia. *PLoS Global Public Health*, 2(9), e0000241. <https://doi.org/10.1371/journal.pgph.0000241>
- Budiarti, M., Maruzy, A., Mujahid, R., Sari, A. N., Jokopriyambodo, W., Widayat, T., & Wahyono, S. (2020). The use of antimalarial plants as traditional treatment in Papua Island, Indonesia. *Heliyon*, 6(12), e05562. <https://doi.org/10.1016/j.heliyon.2020.e05562>
- Debora, J., Rinonce, H. T., Pudjohartono, M. F., Astari, P., Winata, M. G., & Kasim, F. (2018). Prevalensi malaria di Asmat, Papua: Gambaran situasi terkini di daerah endemik tinggi. *Journal of Community Empowerment for Health*, 1(1). <https://doi.org/10.22146/jcoemph.38309>
- Dhewantara, P. W., Ipa, M., & Widawati, M. (2019). Individual and contextual factors predicting self-reported malaria among adults in eastern Indonesia: Findings from Indonesian community-based survey. *Malaria Journal*, 18(1), 1-17. <https://doi.org/10.1186/s12936-019-2758-2>
- Guntur, R. D., Kingsley, J., & Amirul Islam, F. M. (2021). Malaria awareness of adults in high, moderate and low transmission settings: A cross-sectional study in rural East Nusa Tenggara Province, Indonesia. *PLoS ONE*, 16(11 November 2021), 1-18. <https://doi.org/10.1371/journal.pone.0259950>
- Guntur, R. D., Kingsley, J., & Islam, F. M. A. (2021). Epidemiology of malaria in east nusa tenggara province in indonesia: Protocol for a cross-sectional study. *JMIR Research Protocols*, 10(4), 1-12. <https://doi.org/10.2196/23545>
- Hasyim, H., Dale, P., Groneberg, D. A., Kuch, U., & Müller, R. (2019). Social determinants of malaria in an endemic area of Indonesia. *Malaria Journal*, 18(1), 1-11. <https://doi.org/10.1186/s12936-019-2760-8>
- Hasyim, H., Nursafingi, A., Haque, U., Montag, D., Groneberg, D. A., Dhimal, M., Kuch, U., & Müller, R. (2018). Spatial modelling of malaria cases associated with environmental factors in South Sumatra, Indonesia. *Malaria Journal*, 17(1), 1-15. <https://doi.org/10.1186/s12936-018-2230-8>
- Herman, H. (2020). The Relationship of Family Roles and Attitudes in Child Care With Cases of Caput Succedeneum in RSUD Labuang Baji, Makassar City in 2018. *Jurnal Inovasi Penelitian*, 1(2), 49-52. <https://doi.org/10.47492/jip.v1i2.49>
- Inriyanti, A., Hukubun, M. C. Y., Ayomi, M. B., & Tappy, M. (2021). Malaria in the Borderlands of Papua Province of the Republic of Indonesia and Papua New Guinea. 25(3), 5755-5766.
- Ipa, M., Laksono, A. D., Astuti, E. P., Prasetyowati, H., & Hakim, L. (2020). Predictors of malaria incidence in rural eastern Indonesia. *Indian Journal of Forensic Medicine and Toxicology*, 14(4), 3105-3111. <https://doi.org/10.37506/ijfmt.v14i4.12078>
- Ipa, M., Widawati, M., Laksono, A. D., Kusri, I., & Dhewantara, P. W. (2020). Variation of preventive practices and its association with malaria infection in eastern Indonesia: Findings from community-based survey. *PLoS ONE*, 15(5), 1-18. <https://doi.org/10.1371/journal.pone.0232909>
- Istiana, I., Hadi, U., Dachlan, Y. P., & Arwati, H. (2021). Malaria at forest areas in south kalimantan, indonesia: Risk factors and strategies for elimination. *Open Access Macedonian Journal of Medical Sciences*, 9, 1147-1154. <https://doi.org/10.3889/oamjms.2021.7012>
- Kemenkes RI. (2019). Indonesia Health Profile 2019. In *Kementrian Kesehatan Republik Indonesia* (Vol. 42, Issue 4).
- Konaté, D., Diawara, S. I., Touré, M., Diakité, S. A. S., Guindo, A., Traoré, K., Diarra, A., Keita, B., Thiam, S., Keita, M., Sissoko, I., Sogoba, N., Traoré, S. F., Krogtad, D. J., Doumbia, S., & Diakité, M. (2020). Effect of routine seasonal malaria chemoprevention on malaria trends in children under 5 years in Dangassa, Mali. *Malaria Journal*, 19(1), 1-6. <https://doi.org/10.1186/s12936-020-03202-y>
- Mahkota, R., Nurcandra, F., Anggraini, F. D. P., Putri, A. I., & Wispriyono, B. (2020). Risk of agricultural



- pesticide exposure to malaria incidence and anopheles susceptibility at an endemic area in central Java, Indonesia – A case-control study. *Open Access Macedonian Journal of Medical Sciences*, 8(E), 52–59. <https://doi.org/10.3889/oamjms.2020.3024>
- Manangsang, F., Ganing, A., Purba, E. R. V, Rumaseb, E., & Sarwadhamana, R. J. (2021). Analysis of Environmental Risk Factors Against Malaria Events in Kerom Province District Papua. *Journal of Hospital Administration*, 4(2), 37–42.
- Morakinyo, O. M., Balogun, F. M., & Fagbamigbe, A. F. (2018). Housing type and risk of malaria among under-five children in Nigeria: Evidence from the malaria indicator survey. *Malaria Journal*, 17(1), 1–11. <https://doi.org/10.1186/s12936-018-2463-6>
- Olubiyi, Folami, A. (2018). Effect of Nursing Intervention on Knowledge of Malaria Prevention among Mothers of Under-Five Children in Selected Primary Health Care, Mushin Local Government, Lagos. *Texila International Journal of Public Health*, 6(1), 83–92. <https://doi.org/10.21522/tijph.2013.06.01.art008>
- Oluwasogo; Henry; Abdulrasheed; Olawuni; and Olabisi. (2016). Assessment of Mother's Knowledge and Attitude towards Malaria Management among Under Five (5) Years Children in Okemesi-Ekiti, Ekiti-West Local Government, Ekiti State Estrjkl. *Malaria Control & Elimination*, 5(2). <https://doi.org/10.4172/2470-6965.1000142>
- Orhan, I., & Serin, E. K. (2019). Use of Health Technologies by Nurses and Their Thoughts on Technology. *International Journal of Caring Sciences*, 12(1), 416–422.
- Ouédraogo, M., Samadoulougou, S., Rouamba, T., Hien, H., Sawadogo, J. E. M., Tinto, H., Alegana, V. A., Speybroeck, N., & Kirakoya-Samadoulougou, F. (2018). Spatial distribution and determinants of asymptomatic malaria risk among children under 5 years in 24 districts in Burkina Faso. *Malaria Journal*, 17(1), 1–12. <https://doi.org/10.1186/s12936-018-2606-9>
- Oyibo, W., Ntadom, G., Uhomoibhi, P., Oresanya, O., Ogbulafor, N., Ajumobi, O., Okoh, F., Maxwell, K., Ezeiru, S., Nwokolo, E., Amajoh, C., Ezeigwe, N., Audu, M., & Conway, D. (2021). Geographical and temporal variation in reduction of malaria infection among children under 5 years of age throughout Nigeria. *BMJ Global Health*, 6(2). <https://doi.org/10.1136/bmjgh-2020-004250>
- Padonou, G. G., Gbenoudon, J. G., Osse, R., Salako, A., Kpanou, C., Sagbohan, H., Gnanguenon, V., Agbo, F. O., Oussou, O., & Akogbeto, M. C. (2018). Knowledge-Attitudes-Practices about Malaria among Communities in Southern Benin. *International Journal of Public Health Science (IJPHS)*, 7(3), 186. <https://doi.org/10.11591/ijphs.v7i3.14395>
- Patriani, D., Arguni, E., Kenangalem, E., Dini, S., Sugiarto, P., Hasanuddin, A., Lampah, D. A., Douglas, N. M., Anstey, N. M., Simpson, J. A., Price, R. N., & Poespoprodjo, J. R. (2019). Early and late mortality after malaria in young children in Papua, Indonesia. *BMC Infectious Diseases*, 19(1), 922. <https://doi.org/10.1186/s12879-019-4497-y>
- Rahmasari, F. V., Setyonugroho, W., Swarjana, I. K., Arisandi, D., & Kesetyaningsih, T. W. (2021). The association between demographic and attitude factors with the practice of malaria prevention among the rural community in Purworejo district, Indonesia. *Qanun Medika - Medical Journal Faculty of Medicine Muhammadiyah Surabaya*, 5(1), 113. <https://doi.org/10.30651/jqm.v5i1.5416>
- Rizkyansah, G., & Rahayu, E. (2021). Implementation of human development policy in health sector in decentralization perspective. *International Journal of Public Health Science*, 10(2), 348–353. <https://doi.org/10.11591/ijphs.v10i2.20671>
- Roosihermatie, B., Widjiartini, Paramita, A., Nugroho, A., & Suprpto, A. (2017). Malaria self-care in Nimboran subdistrict, Jayapura district, Papua province, Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health*, 48(1), 1–8.
- Saha, A., Sarker, M., Kabir, M., Lu, G., & Müller, O. (2019). Knowledge, attitudes, and practices regarding malaria control among the slash and burn cultivators in Rangamati Hill tracts of Bangladesh. *Malaria Journal*, 18(1), 1–9. <https://doi.org/10.1186/s12936-019-2849-0>
- Sembiring, B. M., & Lubis, F. H. (2019). The Relationship Of Knowledge Level With Action Nurses In Medical Waste Management At Sembiring Hospital, Old Deli. *Jurnal Kesehatan Masyarakat & Gizi (Jkg)*, 1(2), 70–77. <https://doi.org/10.35451/jkg.v1i2.171>
- Sroyer, A. M., Mandowen, S. A., & Reba, F. (2022). Analisis Cluster Penyakit Malaria Provinsi Papua Menggunakan Metode Single Linkage Dan K-Means. *Jurnal Nasional Teknologi Dan Sistem Informasi*, 7(3), 147–154. <https://doi.org/10.25077/teknosi.v7i3.2021.147-154>
- Tizifa, T. A., Kabaghe, A. N., McCann, R. S., van den Berg, H., Van Vugt, M., & Phiri, K. S. (2018). Prevention Efforts for Malaria. *Current Tropical Medicine Reports*, 5(1), 41–50. <https://doi.org/10.1007/s40475-018-0133-y>

- Wanzira, H., Katamba, H., Okullo, A. E., Agaba, B., Kasule, M., & Rubahika, D. (2017). Factors associated with malaria parasitaemia among children under 5 years in Uganda: a secondary data analysis of the 2014 Malaria Indicator Survey dataset. *Malaria Journal*, 16(1), 1-9. <https://doi.org/10.1186/s12936-017-1847-3>
- World Health Organization. (2021). *WHO malaria terminology, 2021 update*. <https://apps.who.int/iris/bitstream/handle/10665/349442/9789240038400-eng.pdf?sequence=1&isAllowed=y%0Ahttps://www.who.int/publications/i/item/9789240038400#.YaQBmw6euvI.twitter>
- Yasuoka, J., Kikuchi, K., Nanishi, K., Ly, P., Thavrin, B., Omatsu, T., & Mizutani, T. (2018). Malaria knowledge, preventive actions, and treatment-seeking behavior among ethnic minorities in Ratanakiri Province, Cambodia: A community-based cross-sectional survey 11 Medical and Health Sciences 1117 Public Health and Health Services. *BMC Public Health*, 18(1), 1-11. <https://doi.org/10.1186/s12889-018-6123-0>